

ANASAC KATANA 325 SC

FUNGICIDE

SUSPENSION CONCENTRATE (SC)

Reg. No. L9465

Registration Holder: ANASAC SOUTH AFRICA PTY., LTD

Country: SOUTH AFRICA

GUARANTEED COMPOSITION:

Active ingredient:

AZOXYSTROBIN 20.0%
TEBUCONAZOLE 12.5%

GENERAL

ANASAC KATANA 325 SC should be applied at the signs of infection and before the 5% infection level is reached. A second spray 28 days after the initial application is recommended for wheat with a high yield potential, particularly in case where the initial application was made early (before first/second node stage) or when other diseases develop later in the season. Apply the second application at the flag leaf growth stage.

RECOMMENDATIONS FOR USE

WHEAT	DOSAGE RATES	REMARKS
<p>ANASAC KATANA 325 SC should be applied at the first signs of infection and before the 5% infection level is reached. A second spray 28 days after the initial application is recommended for wheat with a high yield potential, particularly in cases where the initial application was made early (before first/second node stage) or when other diseases develop later in the season. Apply the second application at the flag leaf growth stage.</p>		
<p>Eye spot (<i>Pseudocercospora herpotrichoides</i>) Powdery mildew (<i>Erysiphe graminis</i>) Speckled leaf blotch (<i>Septoria tritici</i>)</p>	<p><i>Ground Application</i> 500mL in 300L water /ha <i>Aerial Application</i> 500mL in 30L water /ha</p>	<p>Eye spot: Spray at the first to second node growth stage. Powdery mildew: Spray when symptoms are noticed but not later than the flag leaf growth stage. Repeat the application if re-infection occurs. Speckled leaf blotch: Spray before the appearance of the flag leaf.</p>

METHODS OF APPLICATION

Sure equipment and nozzles are in good condition and calibrated for a good distribution and uniform droplet size. Before applying the caliber product with water spray equipment to verify that you are applying the correct dose.

MIXING

Verify the correct calibration of the equipment before starting the application. Put water halfway applicator device tank, add the required amount of the product according to the recommended dose in this label, keeping the agitator running and complete with the remaining volume of water. We recommend adding a non-ionic adjuvant to improve coverage.

COMPATIBILITY: The compatibility of Katana 325 SC with other products has not been tested. Katana 325 SC is not compatible with alkaline products. Compatibility with other products is influenced by the formulation of the products involved and the quality of the spray water. A physical compatibility test should always be carried out prior to application.

PHYTOTOXICITY: When tank mixtures with grass herbicides are applied slight leaf mottling/scorch may occur but new growth will not be affected.

RE-ENTRY TO THE AREA TREATED: Do not enter the treated field until the spray deposit has dried unless wearing protective clothing.

WITHHOLDING PERIOD: Allow the following number of days between the last Katana 325 SC application and harvesting or feeding or grazing: Wheat - 77 days.

WARNINGS FOR USE

BEFORE OPENING THE CONTAINER CAREFULLY READ THE PRODUCT LABEL. NO HANDLING WITH THE PRODUCT DIRECTLY IN HAND AND USE with adequate ventilation. During handling, avoid contact with the skin and mucosa not inhale vapors of the formulated product or work in the overspray. Always wear suitable protective elements: long-sleeved coveralls, gloves neoprene, rubber boots, goggles, waterproof hood and mask filters. After use, remember to wash their hands before eating food, drinking, chewing gum, smoking or using th* toilet, bathe with plenty of soap and water, change of clean clothes, never wash your work clothes and protective equipment together with everyday clothing. Use of this product is exclusively agricultural, apply only the recommended doses, do not spray against the wind or higher winds at 10 km / h. Do not use equipment with no loss and unclog pipes with nozzles or mouth.

FIRST AID TREATMENT

Remove the patient from the source of poisoning and keep him/her quiet and reassured.

In the case of skin contact, wash thoroughly with soap and water.

In the case of eye contact, flush with water for at least 15 minutes. Get medical attention if necessary.

If swallowed, do not make the patient vomit. Call a doctor immediately and make the label available to him.

NOTE TO DOCTOR

Do not allow aspiration of gastric contents if gastric lavage performed. Consider use of a laxative. Treat symptomatically.

ANTIDOTE: It does not have a specific antidote

MEASURES FOR THE PROTECTION OF THE ENVIRONMENT

Do not contaminate lakes, fountains, rivers and other waterways, disable empty containers and deliver to authorized collection center empty agrochemical containers, make triple rinse containers.

ANASAC KATANA 325 SC is Toxic to fish. Prevent contamination of dams and rivers

PRESENTATIONS: ¼L, ½L, 1L, 5L, 10L, 20L, 25L.

Importado y distribuido por:

Titular del registro:

ANASAC SOUTH AFRICA PTY., LTD
Company Reg. No 2009/000919/07
P.O. Box 2859, Parklands 2121
Tel: 011-880-3241

**AGRICULTURAL USE
CATEGORY IV - CAUTION**

**KEEP OUT OF REACH OF CHILDREN, UNINFORMED PERSONS AND ANIMALS.
"IN CASE OF POISONING CALL A DOCTOR IMMEDIATELY AND SHOW THE LABEL, BOOKLET
CONTAINER OR HEALTH PERSONNEL".
"MAKE THE TRIPLE WASH CONTAINERS, DISABLE, AND DELETE AGREEMENT WITH
INSTRUCTIONS OF COMPETENT AUTHORITIES"
STORE IN THE ORIGINAL CONTAINER UNDER LOCK AND KEY AWAY FROM FOOD AND
FEEDSTUFFS. "
"NO RINSE CONTAINERS OR APPLICATION EQUIPMENT LAKES, RIVERS AND OTHER WATER
SOURCES"
"NOT GO BACK TO THE AREA TREATED BEFORE TIME SUITABLE FOR RE-ENTRY**

INSTRUCTIONS FOR USE AND HANDLING

"BEFORE USING THE PRODUCT CONSULT AN AGRONOMIST"

INFORMATION ON LIABILITY

The manufacturer guarantees that the product was formulated under optimal conditions, meeting the quality standards required by each country as well as the concentrations of active ingredients listed on the label and information sheet. The seller guarantees that the product was stored under optimal conditions, however, offers no guarantee for the result of the product, since applications are subject to varying environmental conditions that are not necessarily the same in which the product was tested. Buyer agrees to be responsible for all risks that could cause the use and handling of the product you are purchasing and undertakes to store under the conditions that the label suggests. The user is responsible for the proper handling and application of the product, as well as use on crops and dose, considering residue tolerances and following label recommendations. The active ingredients repeated with the same mode of action can use to generate the development of resistance by the treated organisms, thereby preventing or hindering proper control. Given the above stated is important to consider within a control strategy, products alternate mode of action. The company is not liable for a lack of control given by the development of resistance. If in doubt, consult our technical department or specialist you trust. The active ingredients repeated with the same mode of action can use to generate the development of resistance by the treated organisms, thereby preventing or hindering proper control. Given the above stated is important to consider within a control strategy, products alternate mode of action. The company is not liable for a lack of control given by the development of resistance. If in doubt, consult our technical department or specialist you trust. The active ingredients repeated with the same mode of action can use to generate the development of resistance by the treated organisms, thereby preventing or hindering proper control. Given the above stated is important to consider within a control strategy, products alternate mode of action. The company is not liable for a lack of control given by the development of resistance. If in doubt, consult our technical department or specialist you trust. The company is not liable for a lack of control given by the development of resistance. If in doubt, consult our technical department or specialist you trust. The company is not liable for a lack of control given by the development of resistance. If in doubt, consult our technical department or specialist you trust.